THE COVID-19 VACCINE:

Moderna, Pfizer, Janssen (Johnson & Johnson)

| | Moderna | Pfizer | Janssen (J&J) |
|------------------------------|--|--|--|
| Safe and Effective? | Ø | | © |
| Side Effects | 83.9%-90.5% of participants had short-term pain at the injection site, and 71.9% -81.9% of participants experienced side effects | 87% of participants had short-term pain at the injection site and 77% of participants experienced side effects | 48.6% of participants had short-term pain at the injection site, and 33.2%–38.9% experienced side effects |
| Large Clinical Trial Size | 30,000 participant randomized, double-blind, placebo-controlled trial | 43,000 participant randomized, double-blind, placebo-controlled trial | 44,000 participant randomized, double-blind, placebo-controlled trial |
| Storage and Handling | Stored in freezer, but stable in refrigerator for 30 days | Stored in ultra-cold freezers | Stored in temperatures 36 to 46 degrees Fahrenheit |
| Dosing | (0.5 ml) - two doses, one month (28 days) apart | Adult: (0.3 ml) - 2 doses, 3 weeks (21 days) apart Pediatric: (0.2 ml) - 2 doses, 3 weeks (21 days) apart | (0.5 ml) single dose |
| Age Range | 18 years and older | 5 years and older | 18 years and older |
| Booster Dose | People ages 18 years and older who received a Moderna primary series should get a booster. Some people can also choose to receive a second booster. | People ages 1 2 years and older who received a Pfizer- BioNTech primary series should get a booster. Some people can also choose to receive a second booster. | Everyone ages 18 years and older should get a booster. Some people can also choose to receive a second booster. People who received one dose of J&J/Janssen COVID-19 vaccine who want a booster are encouraged to get an mRNA COVID-19 vaccine (Pfizer-BioNTech or Moderna). |

